Countdown Round

Berkeley mini Math Tournament

16 November 2013

э

You roll 3 dice. What is the probability that the product of the outcomes is a factor of 64?

э

You roll 3 dice. What is the probability that the product of the outcomes is a factor of 64?

The answer is $\frac{1}{8}$.

Suppose that points A, B, C, D lie on a circle in that order. Segments AC and BD intersect at E. Given that AE = 1, AD = 2, and BE = 3, find all possible values for the length of BC.

白 ト く ヨ

Suppose that points A, B, C, D lie on a circle in that order. Segments AC and BD intersect at E. Given that AE = 1, AD = 2, and BE = 3, find all possible values for the length of BC.

The answer is 6.

Nine math students take a 100 point test. The average of their scores is 79. If a tenth student takes the test, what is the minimum score that she needs to get to raise the class average score to 81?

Nine math students take a 100 point test. The average of their scores is 79. If a tenth student takes the test, what is the minimum score that she needs to get to raise the class average score to 81?

The answer is 99.

Find the unit's digit of the expression $2013^1 + 2013^2 + 2013^3 + ... + 2013^{2013}$.

э

___ ▶ <

Find the unit's digit of the expression $2013^1 + 2013^2 + 2013^3 + ... + 2013^{2013}$.

The answer is 3.

Find the integer closest to $\pi^3 + \pi^2 + \pi + 1$.

æ

ъ

Image: A image: A

Find the integer closest to $\pi^3 + \pi^2 + \pi + 1$.

The answer is 45.

- ● ● ●

æ

Three circles with radii 1, 2, and 3 are mutually externally tangent. Find the area of the triangle formed by their centers.

Three circles with radii 1, 2, and 3 are mutually externally tangent. Find the area of the triangle formed by their centers.

The answer is 6.

If A = (2, 6), P = (5, 15) and B = (7, 21), what is the ratio of the lengths AP : PB?

э

If A = (2, 6), P = (5, 15) and B = (7, 21), what is the ratio of the lengths AP : PB?

The answer is 3 : 2.

э

___ ▶ <

How many three digit numbers are there with an odd number of odd digits?

How many three digit numbers are there with an odd number of odd digits?

The answer is 450.

A square and a triangle have equal area. Let P_s be the perimeter of the square, and P_t be the perimeter of the triangle. What is the maximum possible value of $\frac{P_s^2}{P_t^2}$?

A square and a triangle have equal area. Let P_s be the perimeter of the square, and P_t be the perimeter of the triangle. What is the maximum possible value of $\frac{P_s^2}{P_t^2}$?

The answer is $\frac{4\sqrt{3}}{9}$.

m and *n* are positive integers satisfying $m^2 + n^2 = 2000$. What is the maximum possible value of *mn*?

m and *n* are positive integers satisfying $m^2 + n^2 = 2000$. What is the maximum possible value of *mn*?

The answer is 800.

You are given 9 coins of a special currency, of denominations 1 through 9. In how many ways can you distribute the coins into 5 stacks such that each stack has the same value? Assume the stacks are unordered.

You are given 9 coins of a special currency, of denominations 1 through 9. In how many ways can you distribute the coins into 5 stacks such that each stack has the same value? Assume the stacks are unordered.

The answer is 1.

Every minute it is in your mouth, a stick of gum loses 20% of its current flavor. To the nearest minute, how long will the stick of gum have more than half its original flavor?

Every minute it is in your mouth, a stick of gum loses 20% of its current flavor. To the nearest minute, how long will the stick of gum have more than half its original flavor?

The answer is 3.

Billfred painstakingly wrote all the odd numbers from 1 to 199 (inclusive) on the board. Fredbob accidentally erased the first n odd numbers. If the sum of the remaining numbers on the board is 9900, find n.

Billfred painstakingly wrote all the odd numbers from 1 to 199 (inclusive) on the board. Fredbob accidentally erased the first n odd numbers. If the sum of the remaining numbers on the board is 9900, find n.

The answer is 10.

An infinitely large army of soldiers lies along the line y = x + 2 and advances in the positive x-direction at a constant speed of 1 unit per second. If Jacob is at (0,0) and is running away from the soldiers at a constant speed of 2 units per second, locate where he will be in 2 seconds, assuming he takes the route distancing himself farthest from the nearest soldier.

An infinitely large army of soldiers lies along the line y = x + 2 and advances in the positive x-direction at a constant speed of 1 unit per second. If Jacob is at (0,0) and is running away from the soldiers at a constant speed of 2 units per second, locate where he will be in 2 seconds, assuming he takes the route distancing himself farthest from the nearest soldier.

The answer is $(2\sqrt{2}, -2\sqrt{2})$.

What is the sum of all positive integers less than 105 that are relatively prime to 105?

э

What is the sum of all positive integers less than 105 that are relatively prime to 105?

The answer is 2520.

In how many ways can you rearrange the letters in ADJACENTsuch that T is adjacent to at least one A?

In how many ways can you rearrange the letters in ADJACENTsuch that T is adjacent to at least one A?

The answer is 9360.

Suppose
$$f(x) = x^2 + 28x - 2013$$
.

What is the largest positive integer a such that f(a) is not positive?

___ ▶ <

Suppose
$$f(x) = x^2 + 28x - 2013$$
.

What is the largest positive integer a such that f(a) is not positive?

The answer is 33.

A D

Given triangle ABC with $\overline{AB} = 6$, $\overline{BC} = 8$, and $\angle ABC = 90^{\circ}$. If BD is the altitude from B to AC, what is the length of AD?

Given triangle ABC with $\overline{AB} = 6$, $\overline{BC} = 8$, and $\angle ABC = 90^{\circ}$. If BD is the altitude from B to AC, what is the length of AD?

The answer is $\frac{18}{5}$.

Three rectangles with integer side lengths are placed adjacent to each other to create a larger rectangle R. If the sum of the perimeters of the three rectangles is 42, what is the minimal possible area of R?

Three rectangles with integer side lengths are placed adjacent to each other to create a larger rectangle R. If the sum of the perimeters of the three rectangles is 42, what is the minimal possible area of R?

The answer is 18.

Each of the following self-referential logical statements has a truth value. Determine how many of them are true.

- This statement is both true and false.
- Inis statement is either true or false.
- This statement is neither true nor false.
- If this statement is true, then this statement is false.

Each of the following self-referential logical statements has a truth value. Determine how many of them are true.

- This statement is both true and false.
- Inis statement is either true or false.
- This statement is neither true nor false.
- If this statement is true, then this statement is false.

The answer is 1.

How many non-congruent triangles with integer side lengths have perimeter 8?

э

How many non-congruent triangles with integer side lengths have perimeter 8?

The answer is 1.

A two-digit decimal number XY_{10} is reversed to YX_{16} in hexadecimal. Find XY_{10} .

æ

A two-digit decimal number XY_{10} is reversed to YX_{16} in hexadecimal. Find XY_{10} .

The answer is 53.

Bob multiplies a number by 2 to get 100 as the answer. However, instead of multiplying he was supposed to add the 2. What is the correct answer?

Bob multiplies a number by 2 to get 100 as the answer. However, instead of multiplying he was supposed to add the 2. What is the correct answer?

The answer is 52.

Given that the prime factorization of 123123123 is $3^2 \cdot x \cdot y$, with x < y, find y.

э

A∄ ▶ ∢ ∃=

Given that the prime factorization of 123123123 is $3^2 \cdot x \cdot y$, with x < y, find y.

The answer is 333667.

э

Two standard six-sided dice are rolled. If at least one of the dice has a 2 on top, what is the probability that the dice sum to 9?

Two standard six-sided dice are rolled. If at least one of the dice has a 2 on top, what is the probability that the dice sum to 9?

The answer is 0.

100 Berkeley students are taking a math class. If 77 of them are math majors, 43 of them are statistics majors and 2 of them are computer science majors, what is the greatest number of students who are only music majors that could be in the class if no student is a triple major?

100 Berkeley students are taking a math class. If 77 of them are math majors, 43 of them are statistics majors and 2 of them are computer science majors, what is the greatest number of students who are only music majors that could be in the class if no student is a triple major?

The answer is 23.

Evaluate 47 · 89.

Berkeley mini Math Tournament Countdown Round

æ

@▶ ∢ ≣▶

Evaluate 47 · 89.

The answer is 4183.

æ

- ● ● ●

Two trains are approaching each other at 40 mph and 60 mph respectively. If they are currently 70 miles apart, in how many minutes will they meet?

Two trains are approaching each other at 40 mph and 60 mph respectively. If they are currently 70 miles apart, in how many minutes will they meet?

The answer is 42.